This glossary contains definitions of certain technical terms used in this document in connection with us and our business. These may not correspond to standard industry definitions and may not be comparable to similar terms adopted by other companies.

"Abdominal distension"	a condition where substances, such as air (gas) or fluid, accumulate in the abdomen causing its expansion
"AE(s)"	adverse event, which may be mild, moderate, or severe, any untoward medical occurrences in a patient administered a drug or other pharmaceutical product during clinical trials and which do not necessarily have a causal relationship with the treatment
"affinity"	the extent or fraction to which a drug binds to receptors at any given drug concentration or the firmness with which the drug binds to the receptor. Affinity describes the strength of the attraction between two chemicals, or an antigen and an antibody
"age-related disease(s)"	a disease that is most often seen with increasing frequency with increasing senescence
"age-related macular degeneration" or "AMD"	an irreversible medical condition of partial or complete vision loss caused by degenerative lesions of the retinal pigment epithelium and neuronal retina
"AL"	aluminum hydroxide adjuvant, aluminum hydroxide that is used to increase the efficacy or potency of certain vaccines
"anemia"	a condition where the body does not have enough healthy red blood cells
"ANG2"	angiopoietin-2, part of a family of vascular growth factors that play a role in embryonic and postnatal angiogenesis
"anorexia"	an eating disorder characterized by abnormally low body weight, an intense fear of gaining weight, and a distorted perception of weight
"angiogenesis"	the formation and remodeling of new blood vessels and capillaries from growth of pre-existing blood vessels

"antibody-dependent an immune mechanism through which Fc gamma cell-mediated cytotoxicity" receptor-bearing effector cells can kill target cells or "ADCC" expressing tumor- or pathogen-derived antigens on their surface through antibody-binding effect "anti-angiogenic drugs" a group of targeted drugs that act by inhibiting the growth of the small blood vessels on which cancers depend for their growth through inhibition of VEGF, VEGFR, other related growth factors or receptors "antibody-dependent the mechanism by which antibody-bound target cells cell-mediated phagocytosis" activate the Fc gamma receptors on the surface of or "ADCP" phagocytes to induce phagocytosis, resulting in endocytosis and degradation of the target cell through phagosome acidification "antigen" molecule that stimulates an immune response by activating lymphocytes "apoptosis" programmed cell death, a genetically directed process of cell self-destruction that is marked by the fragmentation of nuclear DNA "aspartate aminotransferase" or an enzyme that helps to determine liver function "aspartate transaminase" "assay" an analysis done to determine (i) the presence of a substance and the amount of that substance and (ii) the biological or pharmacological potency of a drug "BLA" biologics license application "bispecific antibody" or "BsAb" an antibody directed at two different targets or two different epitopes on the same target "BOR" best of response, the best response recorded from the start of the study treatment until the end of treatment taking into account any requirement for confirmation "CAF" cancer-associated fibroblast "carcinoma" a malignant tumor of epithelial origin

"CAR-T" Chimeric Antigen Receptor T-Cell Immunotherapy, a

therapy that uses T cells engineered with chimeric

antigen receptors for cancer treatment

"CD3" cluster of differentiation 3, a protein complex and T cell

co-receptor that is involved in activating the cytotoxic T

cell and T helper cells

"CD38" cluster of differentiation 38, a glycoprotein expressed at

low levels on normal healthy tissues, while at high levels on MM and lymphoma cells, functioning either as a

receptor or as an enzyme

"CDMO(s)" contract development and manufacturing organization,

which is a pharmaceutical company that develops and manufactures drugs for other pharmaceutical companies

on a contractual basis

"cell line" a population of cells that descend from a single cell and

contain the same genetic makeup, thereby producing the same proteins. The productivity of a cell line determines the cost of manufacturing, and the quality of a cell line is

directly related to the quality of the relevant biologics

"cell therapy" a therapeutic method where certain types of immune cells

are specifically treated outside the body and then returned into the body to kill tumor/other pathologic cells through their enhanced immune function and recognition

specificity

"cGMP" or "Current Good current Good Manufacturing Practice (GMP) regulations

Manufacturing Practice" enforced by the FDA that provide for systems that assure proper design, monitoring, and control of manufacturing

processes and facilities. Adherence to the cGMP regulations assures the identity, strength, quality, and purity of drug products by requiring that manufacturers of medications adequately control manufacturing operations. This includes establishing strong quality

operations. This includes establishing strong quality management systems, obtaining appropriate quality raw materials, establishing robust operating procedures,

detecting and investigating product quality deviations,

and maintaining reliable testing laboratories

"CH" the constant domain of the heavy chain of an antibody

"CH1" the first constant domain of the heavy chain of an

antibody

"chemotherapy" a category of cancer treatment that uses one or more

anti-cancer chemotherapeutic agents as part of its

standardized regimen

"CHO" Chinese hamster ovary, the ovary of a small rodent called

the Chinese hamster

"CHO cell(s)" Chinese hamster ovary cell(s)

"choroidal neovascularization" the creation of new blood vessels in the choroid layer of

the eye

"CIK(s)" cytokine-induced killer cell(s)

"CL" the constant domain of the light chain of an antibody

"clinical trial" a research study for validating or finding the therapeutic

effects and side effects of test drugs in order to determine

the therapeutic value and safety of such drugs

"Cmax" the highest concentration of a drug in the blood,

cerebrospinal fluid, or target organ after a dose is given

and before a second dose is given

"CMC" chemistry, manufacturing and controls activities in the

development, licensure, manufacturing and ongoing

marketing of pharmaceutical products

"CMO(s)" contract manufacturing organization, a company that

serves other companies in the pharmaceutical industry on a contract basis to provide comprehensive services from

drug development through drug manufacturing

"cohort" a group of patients as part of a clinical study who share

a common characteristic or experience within a defined

period and who are monitored over time

"combination therapy" treatment in which a patient is given two or more drugs

(or other therapeutic agents) for a single disease

"complement-dependent the mechanism by which antibody-coated target cells cytotoxicity" or "CDC" recruit and activate components of the complement cascade, leading to the formation of a complex of attack on membranes on the cell surface and subsequent cell lysis "compound(s)" a substance formed by two or more ingredients in union "COVID-19" coronavirus disease 2019, a disease caused by the SARS-CoV-2 virus which may cause severe acute respiratory syndrome in humans "CR" complete response, which means that all target lesions have disappeared during the course of treatment. For MA treatment, it means complete disappear of ascites for at least four weeks based on CT evaluation. For MPE treatment, it means the volume of pleural effusion of no greater than 50ml for at least four weeks based on CT evaluation "CRO(s)" contract research organization, a company that provides support to the pharmaceutical, biotechnology, and medical device industries in the form of research and development services outsourced on a contractual basis "CRVO" central retinal vein occlusion, a type of retinal vein occlusion (RVO), an occlusion of the main retinal vein posterior to the lamina cribrosa of the optic nerve and is typically caused by thrombosis "CSO(s)" contract sales organization, a company primarily engaged in providing sales representatives to promote and detail pharmaceutical products "CT" computed tomography, a medical-imaging technique used to obtain detailed internal images of the body "cytokine release syndrome" an acute systemic inflammatory syndrome characterized by fever and multiple organ dysfunction that is associated with therapeutic antibodies, CAR-T therapy, haploidentical allogeneic transplantation

"cytokine(s)" a broad and loose category of small proteins that are important in cell signaling, whose release has an effect on

the behavior of cells expressing corresponding receptors

"cytotoxic" toxic to living cells, causing cell damage or death

"dendritic cell(s)" or "DC(s)" cells that constantly sample their surroundings for

pathogens such as viruses and bacteria, detect dangers, and initiate immune responses. Immature dendritic cells have high endocytic activity and a low T-cells activation potential. Contact with a pathogen induces maturation and the expression of certain cell-surface molecules,

greatly enhancing their ability to activate T cells

"DCR" disease control rate, the total proportion of patients who

demonstrate a response to treatment, equal to the sum of complete responses (CR), partial responses (PR) and

stable disease (SD)

"DLT" dose-limiting toxicity, side effects of a drug or other

treatment that are serious enough to prevent an increase

in the dose of that treatment in clinical trial

"DME" diabetic macular edema, a serious eye complication

characterized by abnormal swellings (edema) in the central part of the retina caused by tiny bulges protruding from the vessel walls, leaking, or oozing fluid and blood

into the retina

"DOR" duration of response, the length of time that a tumor

continues to respond to treatment without the cancer

growing or spreading

"drug product" or "DP" a finished dosage form that contains a drug substance,

generally, but not necessarily in association with other

active or inactive ingredients

"drug substance" or "DS" an active ingredient that is intended to furnish

pharmacological activity or other direct effect in the diagnosis, cure, mitigation, treatment, or prevention of disease or to affect the structure or any function of the human body, but does not include intermediates used in

the synthesis of such ingredient

"EC50" half maximal effective concentration, the concentration of a drug, antibody or toxicant which induces a response halfway between the baseline and maximum "electrolyte disorders" a condition where the level of electrolytes in the body is abnormal epithelial-mesenchymal transition, a process by which "EMT" epithelial cells lose their cell polarity and cell-cell adhesion, and gain migratory and invasive properties to become mesenchymal stem cells "EMT-6" an epithelial tumor cell line that was isolated from the breast of a mouse with a mammary tumor "EpCAM" epithelial cell adhesion molecule, a type I transmembrane glycoprotein which plays a role in epithelial carcinogenesis and is involved in various biological functions, such as cell cycle progression, proliferation, differentiation, and migration, and immune evasion "epitope(s)" the specific part of an antigen to which an antibody attaches itself "Fab" antigen-binding fragment, a region on an antibody that binds to antigens, consisting of a light chain and a VH and a CH1 of the heavy chain "FACS" fluorescence activated cell sorter, a specialized type of flow cytometry that uses fluorescent markers to target and isolate cell groups "Fc" crystallizable fragment, which is the tail region of an antibody that interacts with cell surface receptors called Fc receptors and some proteins of the complement system "FcyR" Fc-gamma receptors, a receptor for the Fc region of immunoglobulin "FDA" the Food and Drug Administration of the United States

"fibrosis" a condition where the body's normal healing process goes

unchecked, leading to the formation of permanent scar

tissue

"first-line" with respect to any disease, the first line therapy, which

is the treatment regimen or regimens that are generally accepted by the medical establishment for initial

treatment

"fusion protein" proteins consisting of at least two domains that are

encoded by separate genes

"Fv" the smallest binding unit of an antibody which consists of

a light-and heavy-chain variable domain

"gamma-glutamyl transpeptidase" an enzyme that helps to transfer molecules

"gastric varices" swollen veins in the lining of the stomach that can bleed

and be life-threatening

"GMP" good manufacturing practices, a system for ensuring that

products are consistently produced and controlled according to quality standards, which is designed to minimize the risks involved in any pharmaceutical production that cannot be eliminated through testing the final product. It is also the practice required in order to conform to the guidelines recommended by agencies that control the authorization and licensing of the

manufacture and sale of pharmaceutical products

"GMT(s)" geometric mean titer, a measurement of the potency and

immunogenicity of a vaccine

"Grade" term used to refer to the severity of adverse events, using

Grade 1, Grade 2, Grade 3, etc.

"granzyme" serine proteases released by cytoplasmic granules within

cytotoxic T cells and NK cells

"hACE2" human angiotensin converting enzyme II

"HCC" hepatocellular carcinoma, a type of cancer arising from

hepatocyte malignant transformation

"heavy chain(s)" the large polypeptide subunit of an antibody "hepatomegaly" a condition where the liver becomes enlarged "HER2" human epidermal growth factor receptor 2 "heterodimer(s)" a protein composed of two polypeptide chains differing in the sequence, number, and kind of their amino acid residues "hIgG1" human immunoglobulin G1 "homodimer(s)" a protein composed of two polypeptide chains that are identical in the sequence, number, and kind of their amino acid residues the use of hormone-based drugs to control or treat a "hormone therapy" certain disease or type of disease "hPD-L1" human programmed death ligand 1 (PD-L1) "HPV" human papillomavirus, a deoxyribonucleic acid (DNA) virus that has many types. HPV is an important cause of cervical cancer and is also associated with other types of genital cancer "hypoalbuminemia" a condition where the level of albumin in the blood is low "hypokalemia" a condition where the level of potassium (K+) in the blood serum is abnormally low "hyponatremia" a condition where the concentration of sodium in the blood is abnormally low "hypovolemia" a condition where there is an abnormally low amount of extracellular fluid in the body "IFN-v" interferon gamma, a dimerized soluble pro-inflammatory cytokine "IgG" immunoglobulin G, the most common antibody type found in blood circulation that plays an important role in

antibody-based immunity against invading pathogens

"IgG1" immunoglobulin G1, a subclass of IgG "IL" interleukin, a type of cytokine and signaling molecule in the immune system to provoke an immune response in the body of a human or other animals "IL-2" interleukin-2, a cytokine with essential roles the immune system, primarily via its direct effects on T cells "IMiDs" immunomodulatory drugs, drugs that regulate cellular and humoral immune functions and can enhance immune function, such as Lenalidomide "immune checkpoint inhibitor(s)" a type of drugs that block the immune evasion of tumor cells by certain molecules, which help promote immune responses and allow immune cells to kill cancer cells "immune myocarditis" a condition where the heart muscle becomes inflamed due to an immune response "immunogenicity" the ability of a particular substance, such as an antigen or epitope, to provoke an immune response in the body of a human and other animal. In other words, immunogenicity is the ability to induce humoral and/or cell-mediated immune responses "immunosuppressants" chemical drugs that inhibit abnormal immune responses in the body and are mainly used in clinical practice to treat inflammatory or autoimmune diseases "immunotherapy" a type of therapy that involves the immune system to help the body fight cancer, infection, and other diseases "IND" investigational new drug or investigational new drug application, also known as clinical trial application in China or the United States. "incidence" the frequency of new cases of a disease in a given population over a given period of time "indication" a symptom or particular circumstance that indicates the advisability or necessity of a specific medical treatment or procedure

"inhibitor" a chemical or substance added or applied to another substance to slow down a reaction or to prevent an unwanted chemical change "innovative drugs under clinical drugs other than current treatment methods for MA or development globally that were **MPE** (including paracentesis, intraperitoneal specifically developed for the intrapleural infusions of chemotherapy drugs, antitreatment of MA and MPE" angiogenic drugs, immunosuppressants on top of paracentesis, and diuretics) that are currently under development, including BsAbs, cell therapies, polypeptides, and other proteins "intestinal root obstruction" a blockage that keeps food or liquid from passing through the intestines "intramuscular injection" or the injection or infusion of a substance into a muscle "intramuscular infusion" "intraperitoneal injection" or the injection or infusion of a substance into the "intraperitoneal infusion" peritoneum (body cavity) "intrapleural injection" or the injection or infusion of a substance into the pleural "intrapleural infusion" cavity "intravenous injection" or an injection or infusion of a substance into a vein and "intravenous infusion" directly into the bloodstream "intravitreal injection" or the injection or infusion of a substance into the vitreous "intravitreal infusion" cavity of the eye "in vitro" Latin for "within the glass," studies using components of an organism that have been isolated from their usual biological surroundings, such as microorganisms, cells or biological molecules "in vivo" Latin for "within the living," studies in which the effects of various biological or chemical substances are tested on whole, living organisms as opposed to a partial or dead organism, or those done in vitro "Knobs-into-Holes" or "KIH" a technology which involves engineering the CH3 domains of antibodies to create a "knob" in one of the two heavy chains and a "hole" in the other heavy chain to promote heterodimerization

"laser photocoagulation" a type of laser surgery, which mainly utilizes the photothermal effect of laser on tissues to locally heat up the target tissue after absorbing laser energy, causing the protein in the tissue to denature and solidify "leukocytosis" a condition characterized by an increase in the number of white blood cells in the blood "leukopenia" a condition characterized by a decrease in the number of white blood cells in the blood "light chain(s)" the small polypeptide subunit of an antibody "local therapy" the topical use of drugs to treat lesions "lymphocyte(s)" a subtype of white blood cells, such as T cells, B cells and NK cells "lymphoma" a type of cancer that starts in the lymphatic system and affects white blood cells called lymphocytes "malignant ascites" or "MA" the accumulation of fluid in the peritoneal cavity resulting from the growth of primary or metastatic malignant neoplasms in the peritoneum "malignant pleural effusion" or the collection of fluid in the pleural cavity resulting from "MPE" malignant disease. Malignant pleural effusions often contain free floating malignant cells "macrophage(s)" a type of white blood cell that plays a role to phagocytose antigens, removes dead cells, and stimulates the action of other immune system cells "MAD" maximum administered dose, the highest dose that is safe to be administered to patients "MDSCs" myeloid-derived suppressor cells, a diverse population of immature myeloid cells that have potent immunesuppressive activity "metastatic" in reference to any disease, including cancer, diseaseproducing organisms or of malignant or cancerous cells transferred to other parts of the body by way of the blood or lymphatic vessels or membranous surfaces

"MM" multiple myeloma, a cancer of the plasma cells in the

bone marrow

"monoclonal antibody" or "mAb" an antibody made by identical immune cells that are all

clones of a unique parent cell, in contrast to polyclonal antibodies which are made from hundreds of different

immune cells

"monotherapy" that uses a single drug to treat a disease or

condition

"mOS" median overall survival, the length of time from either

the date of diagnosis or the start of treatment for a disease that half of the patients in a group of patients diagnosed

with the disease are still alive

"MsAb" multi-specific antibody

"MTD" maximum tolerated dose, the highest dose of a drug or

treatment that does not cause unacceptable side effects

"naïve" not having received therapy

"NDA" new drug application, a process required by a regulatory

authority to approve a new drug for sale and marketing

"neutrophils" a type of white blood cell that act as the immune system's

first line of defense

"neovascularization" the natural formation of new blood vessels

"NK cells" natural killer cells, a type of cytotoxic lymphocyte, which

provides rapid responses to virus-infected cell and other intracellular pathogens, and respond to tumor formation

"NOD/SCID mice" non-obese diabetic/severe combined immunodeficient

mice, a brand of immunodeficient laboratory mice

"NRDL" the National Reimbursement Drug List of China

"NSCLC" non-small cell lung cancer

"objective response rate" or the proportion of patients who have a partial or complete "ORR" response to therapy, a partial response is a decrease in the amount of cancer/in the volume of MA in the body, and a complete response is the disappearance of all signs of cancer/MA in the body "OS" overall survival, a length of time that a patient with a specific disease is still alive, used as a measurement of a drug's effectiveness "other proteins" protein drugs other than mAbs, BsAbs, MsAbs, or antibody fusion proteins, which include cytokines, growth factors, or truncated forms of growth factors "PBMC(s)" peripheral blood mononuclear cell(s) "PCD" programmed cell death, a controlled mechanism that eliminates specific cells under developmental or environmental stimuli "PCT" Patent Cooperation Treaty, an international patent law treaty, which provides a unified procedure for filing patent applications to protect inventions in each of its contracting states "PD-1" programmed death 1, an immune checkpoint receptor expressed on T cells, B cells and macrophages. The normal function of PD-1 is to turn off the T cell mediated immune response as part of the process that stops a healthy immune system from attacking other pathogenic cells in the body. When PD-1 on the surface of a T cell attaches to certain proteins on the surface of a normal cell or a cancer cell, the T cell turns off its ability to kill the cell "PD-L1" programmed death ligand 1, which is a protein on the surface of a normal cell or a cancer cell that binds to its receptor, PD-1, on the surface of the T cell that causes the T cell to turn off its ability to kill the cancer cell "perforin" a glycoprotein responsible for pore formation in cell

membranes of target cells

"peritonitis" a condition that occurs when the thin layer of tissue

inside the abdomen, called the peritoneum, becomes

inflamed

"PFS" progression-free survival, the length of time during and

after the treatment of a disease, such as cancer, that a

patient lives with the disease but it does not get worse

"pharmacodynamics" or "PD" the study of how a drug affects an organism, which,

together with pharmacokinetics, influences dosing,

benefit, and adverse effects of the drug

"pharmacokinetics" or "PK" the study of the bodily absorption, distribution,

metabolism, and excretion of drugs, which, together with pharmacodynamics, influences dosing, benefit, and

adverse effects of the drug

"Phase I clinical trial" study in which a drug is introduced into healthy human

subjects or patients with the target disease or condition and tested for safety, dosage tolerance, absorption, metabolism, distribution, excretion, and if possible, to

gain an early indication of its effectiveness

"Phase II clinical trial" study in which a drug is administered to a limited patient

population to identify possible adverse effects and safety risks, to preliminarily evaluate the efficacy of the product for specific targeted diseases, and to determine dosage

tolerance and optimal dosage

"Phase III clinical trial" study in which a drug is administered to an expanded

patient population generally at geographically dispersed clinical trial sites, in well-controlled clinical trials to generate enough data to statistically evaluate the efficacy and safety of the product for approval, to provide

adequate information for the labeling of the product

"PDT"

"Photo-dynamic therapy" or

corresponding light sources to selectively destroy target tissues through photodynamic reactions. It has been

a treatment method that combines photosensitizers and

widely used in the treatment of various surface tumors,

such as multiple myeloma

"pivotal trial" the clinical trial or study to demonstrate clinical efficacy and safety evidence required before submission for drug marketing approval "placebo" any dummy medical treatment administered to the control group in a controlled clinical trial in order that the specific and non-specific effects of the experimental treatment can be distinguished "pleural sclerosis" pathological tissues composed of fibrous connective tissue, appearing grayish white with granulation tissue on the surface caused by pleural thickening due to pleurisy "PIs" protease inhibitor, a class of compounds that inhibit the activity of protein kinases "polypeptide" compounds formed by three or more amino acid molecules connected together by peptide bonds "pre-clinical studies" studies or programs testing a drug on non-human subjects, to gather efficacy, toxicity, pharmacokinetics and safety information and to decide whether the drug is ready for clinical trials "proteinuria" a condition where there is a high level of protein in the urine "PR" partial response, refers to an at least 30% but below 100% decrease in the size of a tumor or in the extent of cancer in the body in response to treatment, according to the Response Evaluation Criteria in Solid Tumors. For MA and MPE treatment, it means at least 50% reduction in the volume of fluid (ascites or pleural effusion) for at least four weeks based on CT evaluation the concentration of serum to reduce the number of "PRNT50" plaques by 50% compared to the serum-free virus, a measure that tells how much antibody is present or how effective it is "PuFS" puncture-free survival, a length of time to first need for

occurs first

therapeutic puncture or death after treatment, whichever

"Q3W" every three weeks

"QA" quality assurance

"QC" quality control

"RBD" receptor-binding domain, a key part of a virus located on

its "spike" domain that allows it to dock to body receptors to gain entry into cells and lead to infection

"recombinant" the combination of genetic materials from more than one

origin, or a method to express native proteins in vitro by

genetic engineering

"refractory" when used in reference to any type of cancer, cancer that

does not respond to treatment. The cancer may be resistant at the beginning of treatment, or it may become

resistant during treatment

"relapsed" when used in reference to any disease, including cancer,

the return of a disease or the signs and symptoms of a disease after a period of improvement. With respect to cancer, the likely relapse occurs because a few of the original cancer cells survived the initial treatment. Sometimes, this occurs because cancer cells spread to other parts of the body and were too small to be detected

during the follow-up immediately after treatment

"RM(s)" rhesus macaque(s)

"RP2D" recommended Phase II dose

"rrMM" relapsed/refractory multiple myeloma

"RVO" retinal vein occlusion, caused by a blockage of the veins

carrying blood away from the retina of the eye, which can lead to macular edema where fluid becomes trapped within and under the retina, leading to rapid and severe

loss of visual acuity

"SAE(s)" serious adverse events, any medical occurrence in human drug trials that at any dose: results in death; is lifethreatening; requires inpatient hospitalization or causes prolongation of existing hospitalization; results in persistent or significant disability/incapacity; may have caused a congenital anomaly/birth defect; or requires intervention to prevent permanent impairment or damage "salt-bridge" a combination of amino acids with opposite charge through hydrogen bonding and ionic bonding "SARS-CoV-2" severe acute respiratory syndrome coronavirus 2, a strain of coronavirus that causes COVID-19 "scFv" single-chain variable fragment, a fusion protein of the variable regions of the heavy (VH) and light chains (VL) of immunoglobulins, connected with a protein linker peptide "SCLC" small cell lung cancer "second-line" with respect to any disease, the therapy or therapies that are tried when the first-line treatments do not work adequately "sdAb" or "single-domain an antibody which only includes heavy chain variable antibody" domains to bind to the antigen. In other words, this antibody only includes heavy chains "SHRM" subretinal hyperreflective material, a hyperreflective material seen on optical coherence tomography and located under the retina and above the retinal pigment epithelium "SINE" selective inhibitors of nuclear export, drugs that block exportin 1, a protein involved in transport from the cell nucleus to the cytoplasm. "SMO(s)" site management organization, an organization that

provides clinical trial-related services

"solid tumor" an abnormal mass of tissue that usually does not contain cysts or liquid areas. Solid tumors may be benign (not cancer), or malignant (cancer). Different types of solid tumors are named for the type of cells that form them "SPR" surface plasmon resonance, a label-free technology that can measure sequential binding events and is a key part of any analytical toolbox, enabling the examination of dual-target specificities in a bispecific antibody within a single assay "S protein" spike glycoprotein, the largest of the four major structural proteins found in coronaviruses "systemic edema" a condition where there is an accumulation of fluid in the body's tissues "symmetric tetravalent BsAb(s)" a tetravalent BsAb with symmetric structure that can target two different targets at the same time and is bivalent to each target "systematic treatment" any type of cancer treatment that has effects throughout the body rather than being applied directly to the cancer "T cell(s)" a lymphocyte of a type produced or processed by the thymus gland and actively participating in the immune response, which plays a central role in cell-mediated immunity "T cell-engaging BsAb(s)" BsAb that binds TAA on cancer cells and targets on T cells with their two arms, thereby engaging effective T cells and tumor cells "t<sub>1/2</sub>" or "half-life" the period of time required for the concentration or amount of a drug in the body to be reduced to exactly one-half of a given concentration or amount of such drug "targeted therapy" a treatment method that utilizes drugs with specific targets to interfere with the growth, division, and spread of cancer cells to achieve the goal of treating tumors "TAM" tumor-associated macrophage traditional Chinese medicine "TCM"

"TCR" T cell antigen recognition receptor "TEAE(s)" treatment emergent adverse events, an event that emerges during treatment, having been absent pretreatment, or worsens relative to the pretreatment state "TGF-B" transforming growth factor-\u03b3, a versatile cytokine usually overexpressed in advanced tumors and related to poor prognoses "TGF-B1" transforming growth factor beta 1, a polypeptide member of the TGF-B superfamily of cytokines "thoracic" the chest or thorax area of the body "thrombocytopenia" a condition characterized by abnormally low levels of platelets in the blood "TKI" tyrosine kinase inhibitors, a pharmaceutical drug that inhibits tyrosine kinases "TME" tumor microenvironment, the ecosystem that surrounds a tumor inside the body which includes immune cells, the extracellular matrix, blood vessels and other cells, like fibroblasts. A tumor and its microenvironment constantly interact and influence each other, either positively or negatively "TNFa" tumor necrosis factor-α, a cell signaling protein (cytokine) involved in systemic inflammation and one of the cytokines that make up the acute phase reaction "toxicity" the degree to which a substance or a mixture of substances can harm humans or animals. It is expressed generally as a dose response "TRAE(s)" treatment-related adverse events "Treg(s)" regulatory T cell, a subpopulation of T cells that modulate the immune system, maintain tolerance to self-antigens, and prevent autoimmune disease "tumor-associated antigen(s)" or an antigen molecule present on tumor cells and normal "TAA(s)" cells which has been widely used for treating tumors

"TV" tumor volume "wAMD" wet age-related macular degeneration (AMD), an irreversible medical condition of partial or complete vision loss caused by degenerative lesions of the retinal pigment epithelium and neuronal retina where abnormal blood vessel growth stimulated by VEGF under the macula causes blood and fluid to seep into the retina "variants of concern" or "VOCs" a category used for variants of SARS-CoV-2, being linked to rapid spread in human populations (epidemiological data) "VEGF" vascular endothelial growth factor, a family of signaling protein critical for the growth of the new vessels and thereby development of cancer cells. VEGF, including VEGF-A, VEGF-B, VEGF-C and VEGF-D, binds to VEGF receptors (VEGFRs), generally, VEGF-A is referred to as VEGF unless otherwise specified "VL" the variable domain of the light chain of an antibody "VH" the variable domain of the heavy chain of an antibody "VHH" also known as a nanobody, is the variable domain of the heavy chain of a single-domain antibody "vitreous" a clear, gel-like substance that fills the space between the

lens and the retina of the eyeball