

Hong Kong Exchanges and Clearing Limited and The Stock Exchange of Hong Kong Limited take no responsibility for the contents of this announcement, make no representation as to its accuracy or completeness and expressly disclaim any liability whatsoever for any loss howsoever arising from or in reliance upon the whole or any part of the contents of this announcement.



Genscript Biotech Corporation

(Incorporated in the Cayman Islands with limited liability)
(Stock code: 1548)

OVERSEAS REGULATORY ANNOUNCEMENT LATE-BREAKING LONG-TERM OVERALL SURVIVAL DATA FROM CARTITUDE-4 STUDY IN MULTIPLE MYELOMA

This announcement is made by the board of directors (the “**Board**”) of Genscript Biotech Corporation (the “**Company**”, together with its subsidiaries, the “**Group**”) pursuant to Rule 13.10B of the Rules Governing the Listing of Securities on The Stock Exchange of Hong Kong Limited.

Legend Biotech Corporation (“**Legend**”), a non-wholly owned subsidiary of the Company, whose shares are listed by way of American Depositary Shares on the Nasdaq Global Select Market in the United States (“**U.S.**”), has filed a Form 6-K with the United States Securities and Exchange Commission (the “**SEC**”) in relation to Legend’s issue of a press release announcing late-breaking long-term data from the Phase 3 CARTITUDE-4 study that shows a single infusion of CARVYKTI® (ciltacabtagene autoleucel) significantly extended overall survival (OS) in patients with relapsed or lenalidomide-refractory multiple myeloma who have received at least one prior line of therapy, including a proteasome inhibitor (PI) and an immunomodulatory agent, reducing the risk of death by 45 percent versus standard therapies of pomalidomide, bortezomib and dexamethasone (PVd) or daratumumab, pomalidomide and dexamethasone (DPd). CARVYKTI® is now the first and only cell therapy to improve overall survival versus standard therapies for patients with lenalidomide-refractory multiple myeloma as early as second line. These results were presented as a late-breaking oral presentation at the 2024 International Myeloma Society (IMS) Annual Meeting (Abstract #OA-65) in Rio de Janeiro, Brazil.

For details, please refer to the attached Form 6-K. The attachment is the full Form 6-K and the press release as published on the SEC’s website available at <https://www.sec.gov/Archives/edgar/data/1801198/000117184324005381/0001171843-24-005381-index.html>.

CAUTIONARY NOTE REGARDING FORWARD-LOOKING STATEMENTS

Statements in this announcement about future expectations, plans, and prospects, as well as any other statements regarding matters that are not historical facts, constitute “forward-looking statements” within the meaning of The U.S. Private Securities Litigation Reform Act of 1995. These statements include, but are not limited to, statements relating to Legend’s strategies and objectives; statements relating to CARVYKTI®, including Legend’s expectations for CARVYKTI® and its therapeutic potential; statements related to the potential results from ongoing studies in the CARTITUDE clinical development program; and the potential

benefits of Legend's product candidates. The words "anticipate," "believe," "continue," "could," "estimate," "expect," "intend," "may," "plan," "potential," "predict," "project," "should," "target," "will," "would" and similar expressions are intended to identify forward-looking statements, although not all forward-looking statements contain these identifying words. Actual results may differ materially from those indicated by such forward-looking statements as a result of various important factors. Legend's expectations could be affected by, among other things, uncertainties involved in the development of new pharmaceutical products; unexpected clinical trial results, including as a result of additional analysis of existing clinical data or unexpected new clinical data; unexpected regulatory actions or delays, including requests for additional safety and/or efficacy data or analysis of data, or government regulation generally; unexpected delays as a result of actions undertaken, or failures to act, by the third party partners of the Group; uncertainties arising from challenges to Legend's patent or other proprietary intellectual property protection, including the uncertainties involved in the U.S. litigation process; government, industry, and general product pricing and other political pressures; as well as the other factors discussed in the "Risk Factors" section of Legend's Annual Report on Form 20-F filed with the SEC on March 19, 2024. Should one or more of these risks or uncertainties materialize, or should underlying assumptions prove incorrect, actual results may vary materially from those described in this announcement as anticipated, believed, estimated or expected. Any forward-looking statements contained in this announcement speak only as of the date of this announcement. The Company and Legend specifically disclaim any obligation to update any forward-looking statement, whether as a result of new information, future events or otherwise.

This announcement has been issued in the English language with a separate Chinese language translation. If there is any inconsistency or ambiguity between the English version and the Chinese version, the English version shall prevail.

By order of the Board
Genscript Biotech Corporation
Jiange Meng
Chairman and Executive Director

Hong Kong, September 29, 2024

*As at the date of this announcement, the executive Directors are Dr. Fangliang Zhang ("**Dr. Frank Zhang**"), Mr. Jiange Meng ("**Mr. Robin Meng**"), Ms. Ye Wang ("**Ms. Sally Wang**") and Dr. Li Zhu; the non-executive Directors are Dr. Luquan Wang ("**Dr. Larry Wang**") and Dr. Ross Allen Grossman; and the independent non-executive Directors are Mr. Zumian Dai ("**Mr. Edward Dai**"), Mr. Jiuan Pan ("**Mr. Ethan Pan**"), Mr. Yiu Leung Andy Cheung, Dr. Chenyang Shi ("**Dr. Victor Shi**") and Dr. Alphonse Galdes.*

**UNITED STATES
SECURITIES AND EXCHANGE COMMISSION
Washington, D.C. 20549**

FORM 6-K

**REPORT OF FOREIGN PRIVATE ISSUER PURSUANT TO RULE 13a-16 OR 15d-16 UNDER THE SECURITIES
EXCHANGE ACT OF 1934**

Date of Report: September 27, 2024

Commission File Number: **001-39307**

Legend Biotech Corporation

(Translation of registrant's name into English)

2101 Cottontail Lane Somerset, New Jersey 08873

(Address of principal executive office)

Indicate by check mark whether the registrant files or will file annual reports under cover of Form 20-F or Form 40-F.
Form 20-F ☒ Form 40-F ☐

Legend Biotech Announced Late-Breaking Long-Term Overall Survival Data from Landmark CARTITUDE-4 Study in Multiple Myeloma

On September 27, 2024, Legend Biotech Corporation (“Legend Biotech”) issued a press release announcing that late-breaking long-term data from the Phase 3 CARTITUDE-4 study that shows a single infusion of CARVYKTI® (ciltacabtagene autoleucel) significantly extended overall survival (OS) in patients with relapsed or lenalidomide-refractory multiple myeloma who have received at least one prior line of therapy, including a proteasome inhibitor (PI) and an immunomodulatory agent, reducing the risk of death by 45 percent versus standard therapies of pomalidomide, bortezomib and dexamethasone (PVD) or daratumumab, pomalidomide and dexamethasone (DPd). CARVYKTI® is now the first and only cell therapy to improve overall survival versus standard therapies for patients with lenalidomide-refractory multiple myeloma as early as second line. These results were presented as a late-breaking oral presentation at the 2024 International Myeloma Society (IMS) Annual Meeting (Abstract #OA-65) in Rio de Janeiro, Brazil. The press release is attached to this Form 6-K as Exhibit 99.1.

This report on Form 6-K, including Exhibit 99.1 (other than the information included under “About Legend Biotech”), is hereby incorporated herein by reference in the registration statements of Legend Biotech on Form F-3 (Nos. 333-278050, 333-257625, and 333-272222) and Form S-8 (No. 333-239478), to the extent not superseded by documents or reports subsequently filed.

EXHIBIT INDEX

Exhibit	Title
99.1	Press Release, dated September 27, 2024

SIGNATURES

Pursuant to the requirements of the Securities Exchange Act of 1934, the registrant has duly caused this report to be signed on its behalf by the undersigned, thereunto duly authorized.

Legend Biotech Corporation

(Registrant)

Date: September 27, 2024

/s/ Ying Huang

Ying Huang, Ph.D.
Chief Executive Officer

CARVYKTI® is the First and Only Cell Therapy to Significantly Extend Overall Survival Compared to Standard of Care in Patients with Multiple Myeloma as Early as Second Line

- *Landmark CARTITUDE-4 study results show CARVYKTI® reduced the risk of death by 45 percent after three-year follow-up*
- *Late-breaking data featured in an oral presentation at the 21st International Myeloma Society Annual Meeting*

SOMERSET, N.J., Sept. 27, 2024 (GLOBE NEWSWIRE) -- Legend Biotech Corporation (NASDAQ: LEGN) (Legend Biotech), a global leader in cell therapy, today announced late-breaking three-year follow-up data from the Phase 3 CARTITUDE-4 study that shows a single infusion of CARVYKTI® (ciltacabtagene autoleucel) significantly extended overall survival (OS) in patients with relapsed or lenalidomide-refractory multiple myeloma who have received at least one prior line of therapy, including a proteasome inhibitor (PI), and an immunomodulatory agent (IMiD), reducing the risk of death by 45 percent versus standard therapies of pomalidomide, bortezomib and dexamethasone (PVD) or daratumumab, pomalidomide and dexamethasone (DPd).¹ CARVYKTI® is now the first and only cell therapy to improve OS versus standard therapies for patients with lenalidomide-refractory multiple myeloma as early as second line.¹ These results were presented as a late-breaking oral presentation at the 2024 International Myeloma Society (IMS) Annual Meeting (Abstract #OA-65) in Rio de Janeiro, Brazil.¹

“These long-term results are groundbreaking and demonstrate that CARVYKTI significantly extends patients’ overall survival and meaningfully improves their quality of life,” said Maria-Victoria Mateos, M.D., Ph.D., Consultant Physician in the Hematology Department and Associate Professor of Medicine at the University of Salamanca, Spain. “These data show that with a single infusion, we can reduce the risk of death and offer multiple myeloma patients the opportunity to live longer.”[‡]

“These data underscore Legend’s commitment to providing patients with effective treatments that have the potential to improve their quality of life and prolong survival,” said Ying Huang, Ph.D., Chief Executive Officer of Legend Biotech. “We will continue to investigate the benefits CAR-T can bring to patients by investing significantly in R&D and exploring the full spectrum of CAR-T technologies.”

The Phase 3 CARTITUDE-4 study evaluated CARVYKTI® compared to standard therapies of PVD or DPd for the treatment of patients with relapsed or lenalidomide-refractory multiple myeloma after one prior line of therapy.¹ Patients who received one to three prior lines of therapy, including a PI and IMiD, and were lenalidomide-refractory were randomized [cilta-cel, n=208, standard therapies, n=211].¹ At median follow-up of almost three years (34 months), median OS was not reached [NR] for patients in the CARVYKTI® arm (95 percent Confidence Interval [CI], not estimable (NE) – NE) or patients that received standard therapies (95 percent CI, 37.75 months – NE) (Hazard Ratio [HR], 0.55; 95 percent CI, 0.39-0.79, $P=0.0009$).¹ At 30-months follow up, OS rates were 76 percent for patients in the CARVYKTI® arm and 64 percent for patients treated with standard therapies.¹ These data show CARVYKTI® significantly extended OS for patients compared to standard therapies.¹

In patients randomized to the CARVYKTI® arm, CARVYKTI® reduced the risk of death by 45 percent compared to standard therapies demonstrating clinically meaningful responses for patients as early as after first relapse.¹ Median progression-free survival (PFS) was NR in the CARVYKTI® arm (95 percent CI, 34.50 months – NE) and 11.79 months (95 percent CI, 9.66-14.00) in patients treated with standard therapies demonstrating sustained deep and durable responses.¹ Patients in the CARVYKTI® arm had a 77 percent complete response or better, and 85 percent overall response rate. Patients in the CARVYKTI® arm had 62

percent minimal residual disease (MRD) negativity at 10^{-5} and 57 percent MRD negativity at 10^{-6} , compared to patients treated with standard therapies (18.5 percent, 9 percent), respectively.¹ Median duration of response was NR (95 percent CI, NE-NE) in the CARVYKTI[®] arm and 18.69 months (95 percent CI, 12.91-23.72) for patients treated with standard therapies.¹ Median time to symptom worsening based on Multiple Myeloma Symptom and Impact Questionnaire (MySIm-Q) was NR (95 percent CI, NE-NE) with CARVYKTI[®] and 34.33 months (95 percent CI, 32.20-NE) with patients treated with standard therapies (HR, 0.38, 95 percent CI, 0.24-0.61; $p < 0.0001$).¹

The safety profile of cilta-cel versus standard therapies was consistent with previous results. In the safety analysis [cilta-cel, n=208, standard therapies, n=208], 97 percent of patients in both arms experienced grade 3/4 treatment-emergent adverse events (TEAEs) with cytopenia being the most common.¹ Treatment-emergent infections occurred in 63 percent of patients in the CARVYKTI[®] arm and 76 percent of patients who received standard therapies with 28 percent and 30 percent being classified as grade 3/4, respectively.¹ In the CARVYKTI[®] arm, there were seven patients with hematologic second primary malignancies, 50 patients died and 21 of those patients died due to progressive disease.¹ Among patients that received standard therapies, one patient experienced a hematologic second primary malignancy, 82 patients died and of those patients, 51 were due to progressive disease.¹

Data from CARTITUDE-4 supported the U.S. Food and Drug Administration (FDA) and European Commission approval of CARVYKTI[®] earlier this year for the treatment of adult patients with relapsed or refractory multiple myeloma who have received at least one prior line of therapy including a PI, IMiD, and are refractory to lenalidomide. Globally, we have now launched CARVYKTI[®] in 5 countries and treated more than 3,500 patients.

CARVYKTI[®] IMPORTANT SAFETY INFORMATION

WARNING: CYTOKINE RELEASE SYNDROME, NEUROLOGIC TOXICITIES, HLH/MAS, PROLONGED and RECURRENT CYTOPENIA, and SECONDARY HEMATOLOGICAL MALIGNANCIES

Cytokine Release Syndrome (CRS), including fatal or life-threatening reactions, occurred in patients following treatment with CARVYKTI[®]. Do not administer CARVYKTI[®] to patients with active infection or inflammatory disorders. Treat severe or life-threatening CRS with tocilizumab or tocilizumab and corticosteroids.

Immune Effector Cell-Associated Neurotoxicity Syndrome (ICANS), which may be fatal or life-threatening, occurred following treatment with CARVYKTI[®], including before CRS onset, concurrently with CRS, after CRS resolution, or in the absence of CRS. Monitor for neurologic events after treatment with CARVYKTI[®]. Provide supportive care and/or corticosteroids as needed.

Parkinsonism and Guillain-Barré syndrome (GBS) and their associated complications resulting in fatal or life-threatening reactions have occurred following treatment with CARVYKTI[®].

Hemophagocytic Lymphohistiocytosis/Macrophage Activation Syndrome (HLH/MAS), including fatal and life-threatening reactions, occurred in patients following treatment with CARVYKTI[®]. HLH/MAS can occur with CRS or neurologic toxicities.

Prolonged and/or recurrent cytopenias with bleeding and infection and requirement for stem cell transplantation for hematopoietic recovery occurred following treatment with CARVYKTI[®].

Secondary hematological malignancies, including myelodysplastic syndrome and acute myeloid leukemia, have occurred in patients following treatment with CARVYKTI®. T-cell malignancies have occurred following treatment of hematologic malignancies with BCMA- and CD19-directed genetically modified autologous T-cell immunotherapies, including CARVYKTI®.

CARVYKTI® is available only through a restricted program under a Risk Evaluation and Mitigation Strategy (REMS) called the CARVYKTI® REMS Program.

WARNINGS AND PRECAUTIONS

INCREASED EARLY MORTALITY - In CARTITUDE-4, a (1:1) randomized controlled trial, there was a numerically higher percentage of early deaths in patients randomized to the CARVYKTI® treatment arm compared to the control arm. Among patients with deaths occurring within the first 10 months from randomization, a greater proportion (29/208; 14%) occurred in the CARVYKTI® arm compared to (25/211; 12%) in the control arm. Of the 29 deaths that occurred in the CARVYKTI® arm within the first 10 months of randomization, 10 deaths occurred prior to CARVYKTI® infusion, and 19 deaths occurred after CARVYKTI® infusion. Of the 10 deaths that occurred prior to CARVYKTI® infusion, all occurred due to disease progression, and none occurred due to adverse events. Of the 19 deaths that occurred after CARVYKTI® infusion, 3 occurred due to disease progression, and 16 occurred due to adverse events. The most common adverse events were due to infection (n=12).

CYTOKINE RELEASE SYNDROME (CRS), including fatal or life-threatening reactions, occurred following treatment with CARVYKTI®. Among patients receiving CARVYKTI® for RRMM in the CARTITUDE-1 & 4 studies (N=285), CRS occurred in 84% (238/285), including ≥ Grade 3 CRS (ASCT 2019) in 4% (11/285) of patients. Median time to onset of CRS, any grade, was 7 days (range: 1 to 23 days). CRS resolved in 82% with a median duration of 4 days (range: 1 to 97 days). The most common manifestations of CRS in all patients combined (≥ 10%) included fever (84%), hypotension (29%) and aspartate aminotransferase increased (11%). Serious events that may be associated with CRS include pyrexia, hemophagocytic lymphohistiocytosis, respiratory failure, disseminated intravascular coagulation, capillary leak syndrome, and supraventricular and ventricular tachycardia. CRS occurred in 78% of patients in CARTITUDE-4 (3% Grade 3 to 4) and in 95% of patients in CARTITUDE-1 (4% Grade 3 to 4). Identify CRS based on clinical presentation. Evaluate for and treat other causes of fever, hypoxia, and hypotension. CRS has been reported to be associated with findings of HLH/MAS, and the physiology of the syndromes may overlap. HLH/MAS is a potentially life-threatening condition. In patients with progressive symptoms of CRS or refractory CRS despite treatment, evaluate for evidence of HLH/MAS. Please see Section 5.4; Hemophagocytic Lymphohistiocytosis (HLH)/Macrophage Activation Syndrome (MAS).

Ensure that a minimum of two doses of tocilizumab are available prior to infusion of CARVYKTI®.

Of the 285 patients who received CARVYKTI® in clinical trials, 53% (150/285) patients received tocilizumab; 35% (100/285) received a single dose, while 18% (50/285) received more than 1 dose of tocilizumab. Overall, 14% (39/285) of patients received at least one dose of corticosteroids for treatment of CRS.

Monitor patients at least daily for 10 days following CARVYKTI® infusion at a REMS-certified healthcare facility for signs and symptoms of CRS. Monitor patients for signs or symptoms of CRS for at least 4 weeks after infusion. At the first sign of CRS, immediately institute treatment with supportive care, tocilizumab, or tocilizumab and corticosteroids.

Counsel patients to seek immediate medical attention should signs or symptoms of CRS occur at any time.

NEUROLOGIC TOXICITIES, which may be severe, life-threatening, or fatal, occurred following treatment with CARVYKTI®. Neurologic toxicities included ICANS, neurologic toxicity with signs and symptoms of parkinsonism, GBS, immune mediated myelitis, peripheral neuropathies, and cranial nerve

palsies. Counsel patients on the signs and symptoms of these neurologic toxicities, and on the delayed nature of onset of some of these toxicities. Instruct patients to seek immediate medical attention for further assessment and management if signs or symptoms of any of these neurologic toxicities occur at any time.

Among patients receiving CARVYKTI® in the CARTITUDE-1 & 4 studies for RRMM, one or more neurologic toxicities occurred in 24% (69/285), including \geq Grade 3 cases in 7% (19/285) of patients. Median time to onset was 10 days (range: 1 to 101) with 63/69 (91%) of cases developing by 30 days. Neurologic toxicities resolved in 72% (50/69) of patients with a median duration to resolution of 23 days (range: 1 to 544). Of patients developing neurotoxicity, 96% (66/69) also developed CRS. Subtypes of neurologic toxicities included ICANS in 13%, peripheral neuropathy in 7%, cranial nerve palsy in 7%, parkinsonism in 3%, and immune mediated myelitis in 0.4% of the patients.

Immune Effector Cell-Associated Neurotoxicity Syndrome (ICANS): Patients receiving CARVYKTI® may experience fatal or life-threatening ICANS following treatment with CARVYKTI®, including before CRS onset, concurrently with CRS, after CRS resolution, or in the absence of CRS.

Among patients receiving CARVYKTI® in the CARTITUDE-1 & 4 studies, ICANS occurred in 13% (36/285), including Grade ≥ 3 in 2% (6/285) of the patients. Median time to onset of ICANS was 8 days (range: 1 to 28 days). ICANS resolved in 30 of 36 (83%) of patients with a median time to resolution of 3 days (range: 1 to 143 days). Median duration of ICANS was 6 days (range: 1 to 1229 days) in all patients including those with ongoing neurologic events at the time of death or data cut-off. Of patients with ICANS, 97% (35/36) had CRS. The onset of ICANS occurred during CRS in 69% of patients, before and after the onset of CRS in 14% of patients respectively.

Immune Effector Cell-associated Neurotoxicity Syndrome (ICANS) occurred in 7% of patients in CARTITUDE-4 (0.5% Grade 3) and in 23% of patients in CARTITUDE-1 (3% Grade 3). The most frequent ≥ 2 % manifestations of ICANS included encephalopathy (12%), aphasia (4%), headache (3%), motor dysfunction (3%), ataxia (2%), and sleep disorder (2%) [see Adverse Reactions (6.1)].

Monitor patients at least daily for 10 days following CARVYKTI® infusion at the REMS-certified healthcare facility for signs and symptoms of ICANS. Rule out other causes of ICANS symptoms. Monitor patients for signs or symptoms of ICANS for at least 4 weeks after infusion and treat promptly. Neurologic toxicity should be managed with supportive care and/or corticosteroids as needed [see Dosage and Administration (2.3)].

Parkinsonism: Neurologic toxicity with parkinsonism has been reported in clinical trials of CARVYKTI®.

Among patients receiving CARVYKTI® in the CARTITUDE-1 & 4 studies, parkinsonism occurred in 3% (8/285), including Grade ≥ 3 in 2% (5/285) of the patients. Median time to onset of parkinsonism was 56 days (range: 14 to 914 days). Parkinsonism resolved in 1 of 8 (13%) of patients with a median time to resolution of 523 days. Median duration of parkinsonism was 243.5 days (range: 62 to 720 days) in all patients including those with ongoing neurologic events at the time of death or data cut-off. The onset of parkinsonism occurred after CRS for all patients and after ICANS for 6 patients.

Parkinsonism occurred in 1% of patients in CARTITUDE-4 (no Grade 3 to 4) and in 6% of patients in CARTITUDE-1 (4% Grade 3 to 4).

Manifestations of parkinsonism included movement disorders, cognitive impairment, and personality changes. Monitor patients for signs and symptoms of parkinsonism that may be delayed in onset and managed with supportive care measures. There is limited efficacy information with medications used for the treatment of Parkinson's disease for the improvement or resolution of parkinsonism symptoms following CARVYKTI® treatment.

Guillain-Barré Syndrome: A fatal outcome following GBS occurred following treatment with CARVYKTI® despite treatment with intravenous immunoglobulins. Symptoms reported include those consistent with Miller-Fisher variant of GBS, encephalopathy, motor weakness, speech disturbances, and polyradiculoneuritis.

Monitor for GBS. Evaluate patients presenting with peripheral neuropathy for GBS. Consider treatment of GBS with supportive care measures and in conjunction with immunoglobulins and plasma exchange, depending on severity of GBS.

Immune Mediated Myelitis: Grade 3 myelitis occurred 25 days following treatment with CARVYKTI® in CARTITUDE-4 in a patient who received CARVYKTI® as subsequent therapy. Symptoms reported included hypoesthesia of the lower extremities and the lower abdomen with impaired sphincter control. Symptoms improved with the use of corticosteroids and intravenous immune globulin. Myelitis was ongoing at the time of death from other cause.

Peripheral Neuropathy: occurred following treatment with CARVYKTI®. Among patients receiving CARVYKTI® in the CARTITUDE-1 & 4 studies, peripheral neuropathy occurred in 7% (21/285), including Grade ≥ 3 in 1% (3/285) of the patients. Median time to onset of peripheral neuropathy was 57 days (range: 1 to 914 days). Peripheral neuropathy resolved in 11 of 21 (52%) of patients with a median time to resolution of 58 days (range: 1 to 215 days). Median duration of peripheral neuropathy was 149.5 days (range: 1 to 692 days) in all patients including those with ongoing neurologic events at the time of death or data cut-off.

Peripheral neuropathies occurred in 7% of patients in CARTITUDE-4 (0.5% Grade 3 to 4) and in 7% of patients in CARTITUDE-1 (2% Grade 3 to 4). Monitor patients for signs and symptoms of peripheral neuropathies. Patients who experience peripheral neuropathy may also experience cranial nerve palsies or GBS.

Cranial Nerve Palsies: occurred following treatment with CARVYKTI®. Among patients receiving CARVYKTI® in the CARTITUDE-1 & 4 studies, cranial nerve palsies occurred in 7% (19/285), including Grade ≥ 3 in 1% (1/285) of the patients. Median time to onset of cranial nerve palsies was 21 days (range: 17 to 101 days). Cranial nerve palsies resolved in 17 of 19 (89%) of patients with a median time to resolution of 66 days (range: 1 to 209 days). Median duration of cranial nerve palsies was 70 days (range: 1 to 262 days) in all patients including those with ongoing neurologic events at the time of death or data cut-off. Cranial nerve palsies occurred in 9% of patients in CARTITUDE-4 (1% Grade 3 to 4) and in 3% of patients in CARTITUDE-1 (1% Grade 3 to 4).

The most frequent cranial nerve affected was the 7th cranial nerve. Additionally, cranial nerves III, V, and VI have been reported to be affected.

Monitor patients for signs and symptoms of cranial nerve palsies. Consider management with systemic corticosteroids, depending on the severity and progression of signs and symptoms.

HEMOPHAGOCYTIC LYMPHOHISTIOCYTOSIS (HLH)/MACROPHAGE ACTIVATION

SYNDROME (MAS): Among patients receiving CARVYKTI® in the CARTITUDE-1 & 4 studies, HLH/MAS occurred in 1% (3/285) of patients. All events of HLH/MAS had onset within 99 days of receiving CARVYKTI®, with a median onset of 10 days (range: 8 to 99 days) and all occurred in the setting of ongoing or worsening CRS. The manifestations of HLH/MAS included hyperferritinemia, hypotension, hypoxia with diffuse alveolar damage, coagulopathy and hemorrhage, cytopenia, and multi-organ dysfunction, including renal dysfunction and respiratory failure.

Patients who develop HLH/MAS have an increased risk of severe bleeding. Monitor hematologic parameters in patients with HLH/MAS and transfuse per institutional guidelines. Fatal cases of HLH/MAS occurred following treatment with CARVYKTI®.

HLH is a life-threatening condition with a high mortality rate if not recognized and treated early. Treatment of HLH/MAS should be administered per institutional standards.

CARVYKTI® REMS: Because of the risk of CRS and neurologic toxicities, CARVYKTI® is available only through a restricted program under a Risk Evaluation and Mitigation Strategy (REMS) called the CARVYKTI® REMS.

Further information is available at <https://www.carvyktirems.com> or 1-844-672-0067.

PROLONGED AND RECURRENT CYTOPENIAS: Patients may exhibit prolonged and recurrent cytopenias following lymphodepleting chemotherapy and CARVYKTI® infusion. Among patients receiving CARVYKTI® in the CARTITUDE-1 & 4 studies, Grade 3 or higher cytopenias not resolved by day 30 following CARVYKTI® infusion occurred in 62% (176/285) of the patients and included thrombocytopenia 33% (94/285), neutropenia 27% (76/285), lymphopenia 24% (67/285) and anemia 2% (6/285). After Day 60 following CARVYKTI® infusion 22%, 20%, 5%, and 6% of patients had a recurrence of Grade 3 or 4 lymphopenia, neutropenia, thrombocytopenia, and anemia respectively, after initial recovery of their Grade 3 or 4 cytopenia. Seventy-seven percent (219/285) of patients had one, two, or three or more recurrences of Grade 3 or 4 cytopenias after initial recovery of Grade 3 or 4 cytopenia. Sixteen and 25 patients had Grade 3 or 4 neutropenia and thrombocytopenia, respectively, at the time of death.

Monitor blood counts prior to and after CARVYKTI® infusion. Manage cytopenias with growth factors and blood product transfusion support according to local institutional guidelines.

INFECTIONS: CARVYKTI® should not be administered to patients with active infection or inflammatory disorders. Severe, life-threatening, or fatal infections, occurred in patients after CARVYKTI® infusion.

Among patients receiving CARVYKTI® in the CARTITUDE-1 & 4 studies, infections occurred in 57% (163/285), including ≥Grade 3 in 24% (69/285) of patients. Grade 3 or 4 infections with an unspecified pathogen occurred in 12%, viral infections in 6%, bacterial infections in 5%, and fungal infections in 1% of patients. Overall, 5% (13/285) of patients had Grade 5 infections, 2.5% of which were due to COVID-19. Patients treated with CARVYKTI® had an increased rate of fatal COVID-19 infections compared to the standard therapy arm.

Monitor patients for signs and symptoms of infection before and after CARVYKTI® infusion and treat patients appropriately. Administer prophylactic, pre-emptive, and/or therapeutic antimicrobials according to the standard institutional guidelines. Febrile neutropenia was observed in 5% of patients after CARVYKTI® infusion and may be concurrent with CRS. In the event of febrile neutropenia, evaluate for infection and manage with broad-spectrum antibiotics, fluids, and other supportive care, as medically indicated. Counsel patients on the importance of prevention measures. Follow institutional guidelines for the vaccination and management of immunocompromised patients with COVID-19.

Viral Reactivation: Hepatitis B virus (HBV) reactivation, in some cases resulting in fulminant hepatitis, hepatic failure, and death, can occur in patients with hypogammaglobulinemia. Perform screening for Cytomegalovirus (CMV), HBV, hepatitis C virus (HCV), human immunodeficiency virus (HIV), or any other infectious agents if clinically indicated in accordance with clinical guidelines before collection of cells for manufacturing. Consider antiviral therapy to prevent viral reactivation per local institutional guidelines/clinical practice.

HYPOGAMMAGLOBULINEMIA: can occur in patients receiving treatment with CARVYKTI®. Among patients receiving CARVYKTI® in the CARTITUDE-1 & 4 studies, hypogammaglobulinemia adverse event was reported in 36% (102/285) of patients; laboratory IgG levels fell below 500mg/dl after infusion in 93% (265/285) of patients.

Hypogammaglobulinemia either as an adverse reaction or laboratory IgG level below 500mg/dl, after infusion occurred in 94% (267/285) of patients treated. Fifty-six percent (161/285) of patients received intravenous immunoglobulin (IVIG) post CARVYKTI® for either an adverse reaction or prophylaxis.

Monitor immunoglobulin levels after treatment with CARVYKTI® and administer IVIG for IgG <400 mg/dL. Manage per local institutional guidelines, including infection precautions and antibiotic or antiviral prophylaxis.

Use of Live Vaccines: The safety of immunization with live viral vaccines during or following CARVYKTI[®] treatment has not been studied. Vaccination with live virus vaccines is not recommended for at least 6 weeks prior to the start of lymphodepleting chemotherapy, during CARVYKTI[®] treatment, and until immune recovery following treatment with CARVYKTI[®].

HYPERSENSITIVITY REACTIONS occurred following treatment with CARVYKTI[®]. Among patients receiving CARVYKTI[®] in the CARTITUDE-1 & 4 studies, hypersensitivity reactions occurred in 5% (13/285), all of which were \leq Grade 2. Manifestations of hypersensitivity reactions included flushing, chest discomfort, tachycardia, wheezing, tremor, burning sensation, non-cardiac chest pain, and pyrexia.

Serious hypersensitivity reactions, including anaphylaxis, may be due to the dimethyl sulfoxide (DMSO) in CARVYKTI[®]. Patients should be carefully monitored for 2 hours after infusion for signs and symptoms of severe reaction. Treat promptly and manage patients appropriately according to the severity of the hypersensitivity reaction.

SECONDARY MALIGNANCIES: Patients treated with CARVYKTI[®] may develop secondary malignancies. Among patients receiving CARVYKTI[®] in the CARTITUDE-1 & 4 studies, myeloid neoplasms occurred in 5% (13/285) of patients (9 cases of myelodysplastic syndrome, 3 cases of acute myeloid leukemia, and 1 case of myelodysplastic syndrome followed by acute myeloid leukemia). The median time to onset of myeloid neoplasms was 447 days (range: 56 to 870 days) after treatment with CARVYKTI[®]. Ten of these 13 patients died following the development of myeloid neoplasms; 2 of the 13 cases of myeloid neoplasm occurred after initiation of subsequent antimyeloma therapy. Cases of myelodysplastic syndrome and acute myeloid leukemia have also been reported in the post-marketing setting. T-cell malignancies have occurred following treatment of hematologic malignancies with BCMA- and CD19-directed genetically modified autologous T-cell immunotherapies, including CARVYKTI[®]. Mature T-cell malignancies, including CAR-positive tumors, may present as soon as weeks following infusions and may include fatal outcomes.

Monitor life-long for secondary malignancies. In the event that a secondary malignancy occurs, contact Janssen Biotech, Inc. at 1-800-526-7736 for reporting and to obtain instructions on collection of patient samples.

EFFECTS ON ABILITY TO DRIVE AND USE MACHINES: Due to the potential for neurologic events, including altered mental status, seizures, neurocognitive decline, or neuropathy, patients receiving CARVYKTI[®] are at risk for altered or decreased consciousness or coordination in the 8 weeks following CARVYKTI[®] infusion. Advise patients to refrain from driving and engaging in hazardous occupations or activities, such as operating heavy or potentially dangerous machinery during this initial period, and in the event of new onset of any neurologic toxicities.

ADVERSE REACTIONS

The most common nonlaboratory adverse reactions (incidence greater than 20%) are pyrexia, cytokine release syndrome, hypogammaglobulinemia, hypotension, musculoskeletal pain, fatigue, infections-pathogen unspecified, cough, chills, diarrhea, nausea, encephalopathy, decreased appetite, upper respiratory tract infection, headache, tachycardia, dizziness, dyspnea, edema, viral infections, coagulopathy, constipation, and vomiting. The most common Grade 3 or 4 laboratory adverse reactions (incidence greater than or equal to 50%) include lymphopenia, neutropenia, white blood cell decreased, thrombocytopenia, and anemia.

Please read full Prescribing Information, including Boxed Warning, for CARVYKTI[®].

ABOUT CARVYKTI[®] (CILTACABTAGENE AUTOLEUCEL; CILTA-CEL)

Ciltacabtagene autoleucel is a BCMA-directed, genetically modified autologous T-cell immunotherapy, which involves reprogramming a patient's own T-cells with a transgene encoding a chimeric antigen receptor (CAR) that identifies and eliminates cells that express BCMA. The cilta-cel CAR protein features two

BCMA-targeting single domain antibodies designed to confer high avidity against human BCMA. Upon binding to BCMA-expressing cells, the CAR promotes T-cell activation, expansion, and elimination of target cells.²

In December 2017, Legend Biotech entered into an exclusive worldwide license and collaboration agreement with Janssen Biotech, Inc. (Janssen), a Johnson & Johnson company, to develop and commercialize cilta-cel. In February 2022, cilta-cel was approved by the U.S. Food and Drug Administration (FDA) under the brand name CARVYKTI[®] for the treatment of adults with relapsed or refractory multiple myeloma. In April 2024, cilta-cel was approved for the second-line treatment of patients with relapsed/refractory myeloma who have received at least one prior line of therapy including a proteasome inhibitor, an immunomodulatory agent, and are refractory to lenalidomide.

In May 2022, the European Commission (EC) granted conditional marketing authorization of CARVYKTI[®] for the treatment of adults with relapsed or refractory multiple myeloma. In September 2022, Japan's Ministry of Health, Labour and Welfare (MHLW) approved CARVYKTI[®]. Cilta-cel was granted Breakthrough Therapy Designation in the U.S. in December 2019 and in China in August 2020. In addition, cilta-cel received a PRiority MEDicines (PRIME) designation from the European Commission in April 2019. Cilta-cel also received Orphan Drug Designation from the U.S. FDA in February 2019, from the European Commission in February 2020, and from the Pharmaceuticals and Medicinal Devices Agency (PMDA) in Japan in June 2020. In March 2022, the European Medicines Agency's Committee for Orphan Medicinal Products recommended by consensus that the orphan designation for cilta-cel be maintained on the basis of clinical data demonstrating improved and sustained complete response rates following treatment.

ABOUT CARTITUDE-4

CARTITUDE-4 (NCT04181827) is an ongoing, international, randomized, open-label Phase 3 study evaluating the efficacy and safety of cilta-cel versus pomalidomide, bortezomib and dexamethasone (PVd) or daratumumab, pomalidomide and dexamethasone (DPd) in adult patients with relapsed or lenalidomide-refractory multiple myeloma who received one to three prior lines of therapy, including a PI and an IMiD.³

ABOUT MULTIPLE MYELOMA

Multiple myeloma is an incurable blood cancer that starts in the bone marrow and is characterized by an excessive proliferation of plasma cells.⁴ In 2024, it is estimated that more than 35,000 people will be diagnosed with multiple myeloma, and more than 12,000 people will die from the disease in the U.S.⁵ While some patients with multiple myeloma initially have no symptoms, most patients are diagnosed due to symptoms that can include bone problems, low blood counts, calcium elevation, kidney problems or infections.⁶

ABOUT LEGEND BIOTECH

Legend Biotech is a global biotechnology company dedicated to treating, and one day curing, life-threatening diseases. Headquartered in Somerset, New Jersey, we are developing advanced cell therapies across a diverse array of technology platforms, including autologous and allogeneic chimeric antigen receptor T-cell, gamma-delta T cell (gd T) and natural killer (NK) cell-based immunotherapy. From our three R&D sites around the world, we apply these innovative technologies to pursue the discovery of cutting-edge therapeutics for patients worldwide.

Learn more at www.legendbiotech.com and follow us on X (formerly Twitter) and LinkedIn.

CAUTIONARY NOTE REGARDING FORWARD-LOOKING STATEMENTS

Statements in this press release about future expectations, plans, and prospects, as well as any other statements regarding matters that are not historical facts, constitute "forward-looking statements" within the meaning of The Private Securities Litigation Reform Act of 1995. These statements include, but are not limited to, statements relating to Legend Biotech's strategies and objectives; statements relating to CARVYKTI[®], including Legend Biotech's expectations for CARVYKTI[®] and its therapeutic potential;

statements related to the potential results from ongoing studies in the CARTITUDE clinical development program; and the potential benefits of Legend Biotech's product candidates. The words "anticipate," "believe," "continue," "could," "estimate," "expect," "intend," "may," "plan," "potential," "predict," "project," "should," "target," "will," "would" and similar expressions are intended to identify forward-looking statements, although not all forward-looking statements contain these identifying words. Actual results may differ materially from those indicated by such forward-looking statements as a result of various important factors. Legend Biotech's expectations could be affected by, among other things, uncertainties involved in the development of new pharmaceutical products; unexpected clinical trial results, including as a result of additional analysis of existing clinical data or unexpected new clinical data; unexpected regulatory actions or delays, including requests for additional safety and/or efficacy data or analysis of data, or government regulation generally; unexpected delays as a result of actions undertaken, or failures to act, by our third party partners; uncertainties arising from challenges to Legend Biotech's patent or other proprietary intellectual property protection, including the uncertainties involved in the U.S. litigation process; government, industry, and general product pricing and other political pressures; as well as the other factors discussed in the "Risk Factors" section of Legend Biotech's Annual Report on Form 20-F filed with the Securities and Exchange Commission on March 19, 2024. Should one or more of these risks or uncertainties materialize, or should underlying assumptions prove incorrect, actual results may vary materially from those described in this press release as anticipated, believed, estimated or expected. Any forward-looking statements contained in this press release speak only as of the date of this press release. Legend Biotech specifically disclaims any obligation to update any forward-looking statement, whether as a result of new information, future events or otherwise

[‡] Maria-Victoria Mateos, M.D., Ph.D., Consultant Physician in the Hematology Department and Associate Professor of Medicine at the University of Salamanca, Spain. has provided consulting, advisory, and speaking services to Legend Biotech; has not been paid for any media work.

INVESTOR CONTACT:

Jessie Yeung
Tel: (732) 956-8271
jessie.yeung@legendbiotech.com

PRESS CONTACT:

Mary Ann Ondish
Tel: (914) 552-4625
media@legendbiotech.com

REFERENCES

¹ Mateos, et al. Overall Survival (OS) With Ciltacabtagene Autoleucel (Cilta-cel) Versus Standard of Care (SoC) in Lenalidomide (Len)-Refractory Multiple Myeloma (MM): Phase 3 CARTITUDE-4 Study Update. International Myeloma Society 2024 Annual Meeting. September 2024.

² CARVYKTI™ Prescribing Information. Horsham, PA: Janssen Biotech, Inc.

³ ClinicalTrials.Gov. A Study Comparing JNJ-68284528, a CAR-T Therapy Directed Against B-cell Maturation Antigen (BCMA), Versus Pomalidomide, Bortezomib and Dexamethasone (PVd) or Daratumumab, Pomalidomide and Dexamethasone (DPd) in Participants With Relapsed and Lenalidomide-Refractory Multiple Myeloma (CARTITUDE-4). <https://www.clinicaltrials.gov/study/NCT04181827>. Accessed March 2024.

⁴ American Cancer Society. "What is Multiple Myeloma?". Available at: <https://www.cancer.org/cancer/types/multiple-myeloma/about/what-is-multiple-myeloma.html>. Accessed March 2024.

⁵ American Cancer Society. "Key Statistics About Multiple Myeloma." Available at: <https://www.cancer.org/cancer/types/multiple-myeloma/about/key-statistics.html>. Accessed March 2024

⁶ American Cancer Society. Multiple myeloma: early detection, diagnosis, and staging. Available at: <https://www.cancer.org/content/dam/CRC/PDF/Public/8740.00.pdf>. Accessed March 2023.